

INDEX.

Abramis cotis (ftnote), 550; gangeticus (ftnote), 550.
 Acanthocheila abducta, 482, 485.
 Acanthochiasma, 177.
 Acanthochiasmida, charac. of, 177.
 Acanthocobitis (ftnote), 539; longipinnis (ftnote), 555.
 Acanthodesmia, 179; skeleton of, 150, 151.
 Acanthodesmida, 173, 175; charac. of, 179.
 Acanthoglossus, 414, 417.
 Acanthometra, 138, 139, 152, 153, 159, 177; tetracopa, 159.
 Acanthometrida, 173, 174, 176.
 Acanthometrina (arctic), 125.
 Acanthonychidæ (*Stimps.*), 647.
 Acanthonychinæ, 647, 648, 649, 650.
 Acanthonyx, charac. of, 650.
 Acanthoplrys, charac. of, 656.
 Acanthophthalmus, 539, 563, 579; pangia, 555.
 Acanthopsis, 539, 563, 579; choirorhynchus, 554; micropogon (ftnote), 555.
 Acanthopterygii, 559, 560.
 Acanthostaurida, charac. of, 177.
 Acanthostaurus, 153, 177.
 Acaridea of Arctic Reg., 121.
 Achæopsis, charac. of, 645.
 Achæus, charac. of, 643.
 Acoura (ftnote), 539; argentata (ftnote), 556; cinerea (ftnote), 555; obscura (ftnote), 556.
 Acourus (ftnote), 539.
 Actinellius, 177.
 Actinolophus pedunculatus, 181.
 Actinomma, 151, 152, 168, 177.
 Actinommatida, 177.
 Actinosphæra, 170.
 Aculeate Hymenoptera, descrip. n. sp. of by F. Smith, 674.
 Adeorbis subcarinatus, 606
 Agelenides, 120.
 Agrypnia pagetana, 727.
 Ailia, 563, 578.
 Ailichthys, 563, 578.
 Ala (*Lockington*), 665.
 Albunea, 326; Gibbesii, 329; Guerini, 326; Lucasii, 330; microps, 328; oxyophthalma, 329; paretii, 330; speciosa, 331; symnista, 326.
 Albuncidæ, 326.
 Alces malchis, anat. of, by Prof. M. Watson and Dr. A. H. Young, 371.
 LINN. JOURN.—ZOOLOGY, VOL. XVI.

Alcyonella fungosa, 490.
 Allman, Prof. G. J., on recent progress in our knowledge of the Structure and Development of Phylactolæmatous Polyzoa, 489; on the Relations of Rhabdopleura, 581.
 Amathia, charac. of, 647.
 Ambassis, 564, 578.
 Amblyceps, 578.
 Amblypharyngodon, 538, 563, 566, 579; Atkinsonii, 544; Jerdoni (ftnote), 544; melittinus, 544; microlepis, 544; mola, 544; pellucidus (ftnote), 544.
 Amblystoma, skull of, 717.
 Ampharete Goëssii, 127.
 Ampharetidæ (arctic), 132.
 Amphicteis Gunneri, 127, 132; Sundevalli, 132, 134.
 Amphictenidæ (arctic), 131.
 Amphilonche, 153, 177.
 Amphipnous, 557, 563, 574, 579; cucchia, 558.
 Amphitrite cirrata, 127.
 Amphiuira filiformis, 447; koreæ, 448, 466, 480; Lütkeni, 448, 464, 480; squamata, 447, 480.
 Amphiuroidæ, 448.
 Amphiuirinae, 448.
 Anabas, 562, 565, 578.
 Anabolia, 730.
 Analysis of the species of Caddis Flies (Phryganea) described by Linnæus in his 'Fauna Suecica.' By H. D. J. Wallengren, with Notes by R. McLachlan, 726.
 Anaptychus, charac. of, 665.
 Anarta Richardsoni, 122.
 Andrenidæ (Sandwich Is.), 682.
 Anematiehthys (ftnote), 538.
 Annelids of British N. Polar Exped., Dr. W. C. McIntosh on, 126.
 Anniversary Address of the President, Prof. Allman, 489.
 Anomalopinae (*Stimps.*), 648.
 Anomalopus (*Stimps.*), nom. præoc., 648.
 Anomalothir, charac. of, 648.
 Anomia, 606; ephippium, 419, 423.
 Anomoura Corystidica, 313; media, 313.
 Anous, 404, 406; cæruleus, 404; cinereus, 404; leucocapillus, 404; melanogenys, 405; stolidus, 405; tenuirostris, 405.

- Anthicus Wollastonii*, 532.
Anthocyrtus, 178.
Anthomyiidae (Arctic reg.), 118.
Antilibinia, charac. of, 650.
Antinoë Sarsi, var. *groenlandica*, 130, 134.
Ants, Fred. Smith on new species from Sandwich Is., 674; Sir J. Lubbock on anatomy of, 738; on habits of, 265, 607.
Ants, affection of, 274; chemical rays of light, effects of on, 285; communications between, 271; exper. with diff. col. lights & media on, 278; exper. with glycerine on, 268; exper. with honey on, 267; exper. testing communication by sound on, 273; kindness of, 624; longevity of, 290, 610; mode of recognition of friends, 610; new species from Sandwich Islands, 675; origin of new communities of, 268; plant-hairs and fur preventative against ants climbing, 607; recollections of friends, 265; relation of second knot and sting of, 622; relations to domestics, 609; smell, 276; sounds emitted by, 623; spectroscopic exper. on, 286; workers fertile as well as queens, but produce males, 607.
Anura, skull of, 718.
Aoria, 247; *Mouloti*, 247.
Aphanias, 535.
Apidae of Arctic region, 106; of Sandwich Is., 684.
Apiomaia, charac. of, 651.
Apis mellifica, 685.
Aplocheilus (fnote), 535; *affinis* (fnote), 535; *carnaticus* (fnote), 535; *chrysostigma* (fnote), 536; *vittatus*, 536.
Appendicularia (arctic), 125.
Aprosmictus chloropterus, 628.
Apterigia (fnote), 557; *immaculata* (fnote), 558; *nigromaculata* (fnote), 558; *saccularis* (fnote), 558.
Apua, 539, 579; *fusca*, 555.
Arachnida of Arctic Region, R. McLachlan on, 98, 120.
Arachnocorys, 178.
Arachnopsis, charac. of, 645.
Arachnosphaeria, 168, 179; *myriacantha*, 152; skeleton of, 150.
Aradasia baccata, 606; *cancellata*, 606.
Aradidae, 485.
Araneida of Arctic reg., 120.
Archebiosis, present state of question, 61.
Arcinella carinata, 421; *laevis*, 422.
Arctic Expedition, Annelids of, Dr. McIntosh on, 126; insects of 78°-83°, R. McLachlan on, 98; surface fauna of, Dr. E. L. Moss on, 122.
Arctopsis, charac. of, 657.
Ardea sacra, 688.
Argynnis chariclea, 109, var. *obscurata*, 110; *polaris*, 109.
Armit, Capt. W. E., on two sp. of *Poëphila*, 97; on *Tachyglossus* & *Ornithorhynchus* in Queensland, 411.
Arses aruensis, 631.
Artamus leucorhynchus, 688.
Aspidomma, 153, 177.
Aspidoparia, 538, 563, 579; *jaya*, 550; *morar*, 550; *sardina* (fnote), 550.
Asteracanthion migratum, 432, 443, 444; *rubens*, 432, 443, 444, var. *migratum*, 432.
Asterioidea of Korean Seas, 424, 443.
Astrolithida, charac. of, 177.
Astrolithium, 153, 177.
Astromma, 176.
Astropecten armatus, 428; *echinulatus*, 426; *euryacanthus*, 428; *formosus*, 424; *hystrix*, 429; *irregularis*, 426, 444; *japonicus*, 427, 443; *Mülleri*, 426; *polyacanthus*, 428, 443, 444; *scoparius*, 428; *vappa*, 428.
Astur torquatus, 627.
Aulacantha, 151, 160, 179.
Aulosphaera, 179; *elegantissima*, 151.
Aulosphaerida, 173, 179.
Australian Finches of the genus *Poëphila*, 95.
Autolytus, 720; *longisetosus*, 131, 134.
Avicula, 606.
Axinus flexuosus, 421; *obesus*, 421, var. *polygona*, 421.
Axonice flexuosa, 132, 134.
Bacilli, their appearance in boiled fluids, 1; in urine, 55.
Bacterium termo, 7, 57.
Badis, 563, 564, 578.
Bagarius, 562, 565, 578.
Balantiocheilus (fnote), 538.
Balitora (fnote), 536.
Baly, Joseph S., Descrip. of new sp. & gen. of *Eumolpidae*, 246; descrip. of n. sp. of *Phytophagous Coleoptera*, 336.
Bangana (fnote), 537.
Barbodes, 575.
Barbus, 538, 561, 566, 574, 579; *ambassis*, 548; *amphibius*, 547; *apogon*, 547; *arenatus*, 547; *arulus*, 546 (fnote) and 547; *Beavani* (fnote), 545; *Blythii*, 546; *bovanicus*, 546; *burmanicus*, 547; *carnaticus*, 545; *caudimarginatus* (fnote), 545; *chagunio*, 545; *chilinoidea*, 545; *chola*, 547; *chrysopoma*, 545; *chrysopterus*, 548; *compressus*, 546; *conchoniis*, 548; *conirostris* (fnote), 548; *cosuatis*,

- 548; *Cumingii*, 548; *curmuca*, 546; *clavatus*, 545; *deliciosus* (fnote), 545; *Denisonii*, 547; *diplochilus* (fnote), 543; *Dobsoni*, 546; *dorsalis*, 547; *dubius*, 545; *dukai*, 545; *Duvaucelli* (fnote), 545; *filamentosus*, 549, and (fnote), 547; *gardinodes* (fnote), 545; *gelius*, 548; *goniosoma*, 545; *gracilis* (fnote), 545; *Grayi* (fnote), 546; *Guentheri* (fnote), 547; *guganio*, 548; *Hamiltonii* (fnote), 546; *hexagonolepis*, 545; *hexastichus*, 546; *himalayanus* (fnote), 545; *innominatus*, 546; *Jerdoni*, 546; *khudree* (fnote), 546; *kolus*, 547; *Layardi* (fnote), 547; *liacanthus* (fnote), 547; *lithopidos*, 546; *McClellandi* (fnote), 548; *macrocephalus* (fnote), 546; *macrolepidotus*, 546; *mahecola*, 547; *mala-baricus*, 546; *megalepis* (fnote), 546; *melanampyx*, 546; *melanostigma*, 547; *micropogon*, 545 (and fnote), 546; *mussulah* (fnote), 546; *mysorensis* (fnote), 545; *nasutus* (fnote), 545; *Neilli*, 546; *nigrofasciatus*, 548; *parral*, 547; *phutunio*, 548; *pleurotania*, 545; *pinnauratus*, 545; *Polydori* (fnote), 545; *progenciis* (fnote), 546; *Puckelli*, 547; *pulchellus*, 546; *punctatus*, 548; *punjaubensis*, 548; *puntio*, 549; *roseipinnis*, 545; *Russelli* (fnote), 545; *sarana*, 545 (and fnote); *sophore*, 546; *spilopholus* (fnote), 545; *spilurus* (fnote), 545; *spinulosus*, 546; *Stephensonii*, 546; *stigma*, 548; *Stoliczkanus*, 548; *Stracheyi*, 546; *subnasutus* (fnote), 545; *terio*, 548; *tetraspilus* (fnote), 547; *tetrarupugus*, 547; *thermalis*, 548; *thomassis*, 546; *tieto*, 548; *tor*, 546; *unimaculatus*, 548; *vittatus*, 549; *Waageni*, 548; *wynaadensis*, 546.
- Barilius*, 538, (fnote) 559, 561, 566, 573, 579; *alburnus* (fnote), 550; *Bakeri*, 551; *barila*, 551; *barna*, 551; *bendelisis*, 551; *Blekeri*, 550; *bola*, 551; *canarensis*, 551; *Evezardi*, 551; *gatensis*, 551; *guttatus*, 551; *morarensis* (fnote), 551; *modestus*, 551; *radiolatus*, 551; *papillatus* (fnote), 551; *rugosus* (fnote), 551; *schaera*, 551; *tileo*, 551; *vagra*, 550.
- Dasilissa* (n. gen.), charac. of, 593; *alta*, 597; *alta*, var. *oxytoma*, 693; *costulata*, 600; *lampra*, 593; *munda*, 596; *oxytropus*, 693; *simplex*, 595; *superba*, 598.
- Bastian*, Prof. H. C., on conditions favouring fermentation and appearance of *Bacilli*, *Micrococci*, and *Torulæ* in Boiled fluids, 1; summary of his views on fermentation and disease, 93.
- Batrachonotus*, charac. of, 645.
- Bdella calandroides*, 121; ?sp. of, 121.
- Bdellidæ*, 121.
- Beale*, Dr., on *Filaria*, 363.
- Bedelia persica*, 259.
- Bees*, Sir J. Lubbock on, 265, 607, 625; species from Sandwich Islands, F. Smith on, 684.
- Belone*, 561, 565, 578; *concola*, 534; *Grayii* (fnote), 534.
- Bembicium*, 603.
- Bembix* (n. gen.), 586; charac. of, 603; *æola*, 603.
- Beroëdes minuta*, 732.
- Beroës* (arctic), 123.
- Bilharzia hæmatobia*, 357.
- Bird-parasites of arctic region*, 118.
- Blackburn*, Rev. Thos., *Hymenoptera of Sandwich Islands* coll. by, 674.
- Blastus* (Leach), 657.
- Blepharopoda*, 333; *occidentalis*, 334; *spinimana*, 335; *spinosa*, 335.
- Bola* (fnote), 538.
- Boletia radiata*, 435.
- Bombus balteatus*, 106; *polaris*, 106.
- Bombyces of arctic region*, 112.
- Bombyx pavonia*, 727; *quereus*, 727.
- Botia*, 539, 563, 579; *almorhæ*, 554; *Berdmorei*, 554; *dario*, 554; *geto*, 554; *histrionica*, 554; *nebulosa*, 554.
- Botryocampe*, 151, 170, 178.
- Botryocytis*, 178.
- Brachelytra of arctic region*, 107.
- Brachiata*, charac. of, 179.
- Brachiopoda*, 418.
- Brachygramma* (fnote), 538.
- Brada villosa*, 127.
- Branched Syllis*, remarkable form of, Dr. W. C. McIntosh on, 720.
- Busk*, George, on recent species of *Heteropora*, 724.
- Butler*, A. G., on *Butterflies in coll. B. M. of gen. Euplœa*, 290.
- Cachius* (fnote), 538.
- Caddis Flies* (*Phryganea*), H. D. J. Wallengren and R. M. Lachlan on, 726.
- Cadulus ampullaceus*, 529; *colubridens*, 523; *curtus*, 527, var. *congruens*, 527; *exiguus*, 528; *gracilis*, 525, 526; *Jeffreysi*, 526; *obesus*, 527; *ovulum*, 529; *rastridens*, 525; *sau-*

- ridens, 525; simillimus, 526; tumidosus, 528; ventricosus, 525; vulpidens, 524.
- Calcaromma calcarea (fntote), 137, 144.
- Callichrous, 562, 565, 578.
- Calliplœa, 295; darchia, 296; dehaanii, 295; doryca 295; hyems, 296; infantilis, 296; Jamesii, 296; ledereri, 295; mazares, 295; niveata, 296; pollita, 296; priapus, 296; pumila, 295; Saundersii, 296; seriata, 295; tulliolus, 296; Turneri, 296; Trimeni, 295.
- Calodietyon, 193.
- Calornis viridescens, 687.
- Camponotus sexguttatus, 675.
- Campophaga spilorrhœa, 688.
- Camposcia, charac. of, 644.
- Cancer carabus (fntote), 327.
- Canceroidea, Dana's separation of (fntote), 641.
- Cantharidus iris, 605.
- Canthophrys albescens (fntote), 554; olivaceus (fntote), 554; rubiginosus (fntote), 555; vittatus (fntote), 554.
- Capitella capitata, 131, 134.
- Capœta (fntote), 537, 538, 575; macrolepidota (fntote), 546; micracanthus (fntote), 540.
- Carpocanium, 178.
- Carpophaga Muellieri, 688; pinon, 688; puella, 688.
- Cataphracta, group of Radiolaria restored, 174; charac. of, 177.
- Catenula, 720.
- Catla, 537, 563, 579; Buchanani, 544.
- Caunopora, 211, 219; (Stromatopora) perforata, 211, 219; placenta, 189, 211, 219; planulata, 210; tubes of (fntote), 212.
- Cellepora mammillata, 238.
- Centropus nigricans, 629.
- Cephus macrocephalus, 689.
- Ceratium divergens, 126; tripos, 125.
- Ceratocarcinus, charac. of, 670.
- Ceratospyris, 178.
- Cervide (anat.) compared with Elk, 388.
- Cervus alces, 371; cashmerianus, 389; elaphus, 388.
- Chaca, 562, 578.
- Chætogaster, 720.
- Chalcopsittacus scintillatus, 686.
- Chalcosoma atlas, 135.
- Chalk, how acted on by limpets, 406.
- 'Challenger' expedition, Mollusca of, Pt. I., 506; Pt. II., 502; Pt. III., 586; Pt. IV., 692.
- Channa, 562, 565, 568, 571.
- Chedrus (fntote), 538; Grayi (fntote), 551.
- Cheilobarbus (fntote), 538.
- Cheiridea (n. gen.), charac. 253; Chaupuisi, 253.
- Chela, 538, 563, 566, 579; alburna (fntote), 552; anastoma (fntote), 552; argentea, 553; bacaila, 553; balooke (fntote), 553; boopis, 553; chipeoides, 553; gora, 553; Owenii (fntote), 553; phulo, 553; punjabensis, 553; sardinella, 553; Sladoni, 553; teekaree (fntote), 553; untrahi, 553.
- Chibia carbonaria, 687.
- Chilomma, 177.
- Chionœetes, charac. of, 654.
- Chironomus, sp. (arctic reg.), 117.
- Chlamydodera cerviniventris, 687.
- Chlamys chinensis, 345; fulvipes, 345; placida, 344; velutina, 343.
- Choleva falklandica, 531.
- Chondrostoma gangeticum (fntote), 543; mullya (fntote), 540; semivelatus (fntote), 542; wattanah (fntote), 543.
- Chone Duneri, 127; infundibuliformis, 133, 134.
- Chorilia, charac. of, 658.
- Chorilibinia, charac. of, 654.
- Chorinus, charac. of, 647.
- Chrysophanus phleas, var. Feildeni, 111.
- Cinclosoma ajax, 631.
- Cirrha, 537, 562, 566, (fntote) 574, 579; bengalensis (fntote), 543; Blochii (fntote), 543; cirrhosa, 543; Dussumieri (fntote), 543; fulungee, 543; latia, 543; macrops (fntote), 543; micropogon (fntote), 541; mrigala, 543; plumbea (fntote), 543; reba, 543; rewah (fntote), 543; rubripinnis (fntote), 543.
- Cirrhinus affinis (fntote), 541; Cuvierii (fntote), 541; macronotus (fntote), 541; rubro-punctatus (fntote), 541.
- Cistenides granulata, 131, 134.
- Cladococcida, 173, 174, 175, 176.
- Cladococcus, 152, 168, 177.
- Cladospyris, 178.
- Clarias, 561, 565, 574, 578; Dussumieri, 571; magur, 571; Teysmanni, 571.
- Classification of Maioid Crustacea or Oxyrhyncha with synop. of Fam., Subfam., and Genera, by Edw. J. Miers, 634.
- Clathrocanium, 178.

- Clathrodictyon, 220; cellulosum, 207, 221, 227; vesiculosum, 220.
 Clio borealis, 124.
 Clitellio arenarius, 133, 134.
 Clupea cyprinoides (fnote), 553; ilisha, 557; sinura (fnote), 557.
 Clypeaster rotundus, 440.
 Clypeastroid sp., 440.
 Cobbold, Prof. T. S., on the life history of *Filaria Bancrofti*, 356, 358.
 Cobitis amnicola (fnote), 554; arenata (fnote), 556; balgara (fnote), 554; bilturio (fnote), 555; bimucronata (fnote), 555; carnaticus (fnote), 554; cinnamomea (fnote), 555; cucura (fnote), 554; gibbosa (fnote), 556; Griffithii (fnote), 556; maya (fnote), 554; microps (fnote), 556; montanus (fnote), 556; moreh (fnote), 555; mysorensis (fnote), 554; ocellata (fnote), 555; oculata (fnote), 554; phoxocheila (fnote), 554; rubripinnis (fnote), 554; scaturigina (fnote), 555; semizonata (fnote), 555; tenuicauda (fnote), 556; vittata (fnote), 556.
 Coccodiscida, charac. of, 176.
 Coccodiscus, 176.
 Cœlocerus, charac. of, 655.
 Cœlodendrida, 173, 174, 175, 176; charac. of, 177.
 Cœlodendrum, 151, 152, 160, 170, 177.
 Cœnostroma, 191; discoideum, 191.
 Colasposoma sellatum, 254.
 Coleoptera (Arctic Region), 102, 107; new, of geographical interest, 530; of Java, 135; new sp. of Phytophagous, 336.
 Colias Boothii, 108; Hecla, var. glacialis, 108.
 Collembola (Arctic Reg.), 119.
 Collida, 157, 173, 174; charac. of, 179.
 Collingwood, C., on n. sp. of Nudi-branches from Eastern Seas, 737.
 Collodes, charac. of, 645.
 Collodinae (*Stimps.*), 644.
 Collonia marginata, 716; munda, 715.
 Collosphæra, 159, 160, 161, 163, 166, 168, 179; Huxleyi, zoospores of, 164.
 Collosphærida, 173.
 Collozoa, group of Radiolaria, 174, 175.
 Collozoum, 159, 160, 161, 162, 166, 179; inerme, 163, 164, 165; colonies of, 138; macrospores of, 165; zoospores of, 164.
 Colpocephalum, sp.? 119; subæquale, 119.
 Conchifera, 419.
 Contributions to Ornithology of New Guinea, R. B. Sharpe's. Pt. V., 626; Pt. VI., 685.
 Copepoda of Arctic Seas, 123, 125.
 Corals compared with Stromatopora, 232.
 Coreidæ, 482.
 Cornutella, 178; scalaris, 148, 150.
 Corvus orru, 687.
 Corystoidea, 313.
 Coscinopora placenta, 189; ramosa, 189.
 Crabro affinis, 677; denticornis, 678; mandibularis, 677; unicolor, 678.
 Crabronidæ (Sandwich Is.), 677.
 Cracticus cassicus, 687; mentalis, 687.
 Crastia, 297; æthiops, 299; aglidice, 298; alecto, 298; Bremeri, 298; camaralzaman, 298; climena, 298; confusa, 299; Crameri, 297; cratis, 297; dioeletia, 297; ebenina, 298; funerea, 298; Goudotii, 298; Grayi, 299; Kinbergi, 297; Lapeyrousei, 299; lugens, 298; malayica, 297; melancholica, 298; melina, 299; modesta, 298; Moorei, 298; nox, 298; occulta, 299; Ochsenheimeri, 297; resarta, 298; Scudderii, 297; sepulchralis, 298; squalida, 298; Wallacei, 299; Zinkenii, 298.
 Crematogaster scutellaris, 283, 290.
 Crenella decussata, 419, 423.
 Crevaux, Dr. J., on "vers de la Guadeloupe," 359.
 Cribrella densispina, 432, 443, 444; sanguinolenta, 444.
 Criocarcinus, charac. of, 661.
 Cristatella, 497.
 Cromyomma, 177.
 Crossochilus (fnote), 537; rostratus (fnote), 543.
 Crustacea, Maioid, or Oxyrhyncha, E. J. Miers on, 634.
 Cryptophagus acutangulus (fnote), 170; gracilipes, 530.
 Cryptopodia, charac. of, 669.
 Cryptus arcticus, 107.
 Culex from Hayes Sound, 116; ? C. caspius, C. nigriceps, or C. pipiens, 117.
 Cyclax, charac. of, 660.
 Cyclocheilichthys (fnote), 538.
 Cyclomaia, charac. of, 660.
 Cyclonox, charac. of, 649.
 Cyclopsitta suavissima, 685.
 Cyclostrema, 700.
 Cynops pyrogaster, skull of, 718.
 Cyphocarcinus, charac. of, 664.
 Cyprinidæ, 559, 560.

- Cyprinodon* (fnote), 559, 561, 578; dispar, 535; *hammonia* (fnote), 535; *lunatus* (fnote), 535; *Stoliczkanus* (fnote), 535.
- Cyprinodontidae*, 535, 559, 560.
- Cyprinus abramioides* (fnote), 544; *acra* (fnote), 542; *apiatus* (fnote), 551; *anjana* (fnote), 549; *cachius* (fnote), 552; *canius* (fnote), 548; *chapalio* (fnote), 552; *chedra* (fnote), 551; *chedrio* (fnote), 551; *cocsa* (fnote), 551; *cura* (fnote), 542; *curechius* (fnote), 541; *cursa* (fnote), 541; *cursis* (fnote), 541; *dancena* (fnote), 553; *dero* (fnote), 542; *falcata* (fnote), 542; *falcatus* (fnote), 542; *goha* (fnote), 551; *gohama* (fnote), 543; *gotyla* (fnote), 540; *Hamiltonii* (fnote), 542; *jogia* (fnote), 549; *kadoon* (fnote), 545; *kakoo* (fnote), 545; *kunnamoo* (fnote), 545; *M'Clellandi* (fnote), 545; *moralis* (fnote), 542; *mosal* (fnote), 546; *musiha* (fnote), 542; *nancar* (fnote), 541; *pausius* (fnote), 542; *putitora* (fnote), 546; *rashora* (fnote), 549; *sada* (fnote), 543; *sophore* (fnote), 548; *sucatio* (fnote), 540; *sutiha* (fnote), 549; *tictis* (fnote), 547; *tila* (fnote), 551; *titius* (fnote), 547.
- Cyrtida*, 141, 157, 173; charac. of, 177.
- Cyrtidosphæra*, 168, 179; skeleton of, 150.
- Cyrtocalpis*, 178; *obliqua*, skeleton of, 150.
- Dacelo Gaudichaudi*, 686; *intermedius*, 686.
- Damæus* near *geniculatus*, 121.
- Danaida* of Latreille, 291.
- Danais similis* as type of *Euplœa*, 291.
- Dangila* (fnote), 537, 575, 579; *Berdmorei*, 543; *burmanica*, 543.
- Danio*, 538, 563, 566, 579; *æquipinnatus*, 552; *albolineata*, 552; *chrysops*, 552; *dangila*, 552; *devario*, 552; *lineatus* (fnote), 552; *lineolatus* (fnote), 552; *malabaricus*, 552; *miconema* (fnote), 552; *neilgherriensis*, 552; *nigrofasciatus*, 552; *rerio*, 552; *spinosus*, 552; *Stoliczkae* (fnote), 552.
- Dascyllus fulvulus*, 135.
- Dasychira grœnlandica*, 112, 117.
- Day, Francis, Geograph. distrib. of Indian Freshwater Fishes, Part III., 534; on occurrence of *Morrhua macrocephala* at mouth of Thames, 698.
- Decipula ovata*, 422.
- Dehaan, remarks on, 650.
- Dendrochelidon mystaceus*, 629.
- Dendronotus* (Arctic), 124.
- Dentalium abyssorum*, 515; *acutissimum*, 514; *agile*, 513; *ægeum*, 509; *amphialum*, 510; *antillarum*, 514; *capillosum*, 508, var. *paucicostatum*, 509; *ceras*, 510; *circumcinctum*, 513; *clavatum*, 524; *compressum*, 516; *dentalis*, 517, 518; *diarrhox*, 511; *didymum*, 517; *ensiculus*, 518; *entalis*, 512, var. *agile*, 512, var. *orthrum*, 512, var. *striolatum*, 512; *erectum*, 513, 514; *fusticulus*, 512; *gadus*, 524; *grande*, 510; *incertum*, 512; *inversum*, 514; *Lamarckii*, 516; *leptosceles*, 512, 513; *Lessoni*, 514; *longitrorsum*, 515; *lubricatum*, 512; *octogonum*, 517; *politum*, 516; *pretiosum*, 515; *semipolitum*, 514; *splendidum*, 514; *subterfissum*, 515, 516; *tornatum*, 518; *yokohamense*, 517; *zelandicum*, 510.
- Dermaleichus stercorarius*, 122.
- Descriptions of new Coleoptera of Geographical interest, by Frederick H. Waterhouse, 530.
- new Hemiptera, by Dr. F. Buchanan White, 482.
- new species Aculeate Hymenoptera from Sandwich Islands, by Fred. Smith, 674.
- new species and genera of Eumolpidae, by J. S. Baly, 246.
- new species of Phytophagous Coleoptera, by J. Baly, 336.
- Deuterocampta Saundersi*, 356.
- Devario* (fnote), 538; *cyanotænia*, (fnote) 552; *M'Clellandi* (fnote), 552.
- Develop. of *Filaria sanguinis hominis*, Dr. P. Manson on, 304.
- of skull of Urodelous Amphibia, W. K. Parker on, 717.
- Dianthia flavicauda* (fnote), 554; *zebra* (fnote), 554.
- Dianthus* (fnote), 539.
- Diaspis Batesi*, 342.
- Diatomacæ* (Arctic), 125.
- Dictyocephalus*, 178.
- Dictyoceras*, 178.
- Dictyocha*, 179; of Baffin's Bay, 125; skeleton of, 150, 151.
- Dictyophimus*, 178.
- Dictyoplegma*, 168.
- Dictyopodium*, 178; sp.? (Wy. Thoms.), 149, 150, 151.
- Dictyosoma*, 155.
- Dictyosphagma*, 155.

- Dictyospyris*, 178.
Dictyostroma, 193, 224; *undulata*, 224.
Dicyrtida, charac. of, 177.
Didymocyrtis, 168, 177.
Dinophysis norvegiana, 126.
Dione (*De Haan*, nom. præoc.), 660.
Diplocheilichthys (fntote), 537.
Diplocheilus (fntote), 537.
Diploconida, 173, 177.
Diploconus, 153, 178; as section of *Radiolaria*, 174.
Diplophysa (fntote), 539.
Diplosphæra, 168, 170, 179.
Diptera of Arctic Region, 104, 116; chitinous integuments of, in salmon's stomach 82° 40' N. lat., 118.
Diptychus, 537, 579; *maculatus*, 541; *Sewerzowi* (fntote), 541.
Dischides bifissus, 521.
Discida, 151, 157, 170, 173; charac. of, 175.
Discognathichthys (fntote), 536.
Discognathus, 536, 560, 561, 566, 574, 579; *Jerdoni*, 540; *lamta*, 540; *macrochir* (fntote), 540; *modestus*, 540.
Discospira, 176.
Discospirida, charac. of, 176.
Disease, origin de novo, 91.
Doclea, charac. of, 652.
Docphorus cebilebrachys, 118.
— sp. on *Bernicla brenta*, 118.
— sp. on *Tetrao rupestris*, 118.
Donacola nigriceps, 688.
Dorataspis, 153, 170, 178; *polyancistra*, 169.
Dorthesia chiton ♀ (fntote), 118.
Doryphora Buckleyi, 352; *dorsomaculata*, 353; *Jacobyi*, 354; *mæsta*, 354; *undulata*, 353; *verrucosa*, 351.
Drymædus Beccarii, 633.
Duncan, Prof. P. M., on *Ophiuroidea* from Korean Seas, 445.
Echidna (*Acanthoglossus*) *Bruijnii*, 414; *hystrix*, 413, 414; in N. & N.E. Queensland, 411; (*Tachyglossus*) *Lawesii*, 414; palate of, 415; *setosa*, 415; skulls of, 413, 415, 416, 417.
Echinanthus testudinarius, 440, 443, 444.
Echinocardium australe, 441, 444; *cor datum*, 442.
Echinoidea of Korean Seas, 424, 443.
Echinolampas oviformis, 441, 443, 444.
Echinometra luenter, 435, 443, 444.
Eclectus polychlorus, 686.
Ectolithia, 173.
Edoliisoma nigrum, 630.
Egeria, charac. of, 654.
Eleotris (fntote), 559, 561, 565, 578.
Elk (*Alces malchis*), anat. of, by Prof. M. Watson and Dr. A. H. Young, 371; cheek of, 372; fore limb, 377; hind limb, 382; kidney of, 374; large intest. of, 373; larynx & circulatory organs of, 374; liver of, 373; myology of, 377; organs of digestion, 371; remarks as to position, 387; small intest. of, 373; spleen of, 373; testicle, &c., of, 374; tongue of, 371; urinary and generative organs of, 374.
Elmis brunneus, 532; *cupreus*, 532.
Emarginula, 606.
Embolus rostralis, 423.
Endocyst of *Polyzoa*, structure of, 491.
Enteromius (fntote), 538.
Entolithia, 173.
Entosphærida as a group, 175; charac. of, 176.
Eolis salmonacea (fntote), 124.
Eos fuscata, 628, 686.
Eozoön, 193.
Epialtus, charac. of, 650; and as syngen. 650.
Erethistes, 563, 578.
Erichoplatus, charac. of, 647.
Erigone provocans, 120; *psychophila*, 120; *vexatrix*, 120.
Eschara nobilis, 237; (*Hemeschara*, *Busk*), 238.
Esomus (fntote), 538; *madraspatensis* (fntote), 549; *malabaricus* (fntote), 549; *vittatus* (fntote), 549.
Esopus, charac. of, 651.
Ethmosphæra, 168, 179.
Ethmosphærida, 173; charac. of, 179.
Etroplus (fntote), 559, 563, 565, 568, 578.
Eubrachus, charac. of, 248; *cylindrica* (type), 248; *indica*, 249; *spinipes*, 248.
Eucecryphalus, 151, 178; *Schultzei*, 140, 150.
Euchelus, 605.
Euchitonia, 153, 176.
Euchone analis, 133, 134; *tuberculosa*, 127.
Eucinetops, charac. of, 644.
Eucrante villosa, 127.
Eucyrtidium, 170, 178; skeleton of, 150, 151.
Euglyptosternum, 573, 578.
Eulabes Dumonti, 687.
Eumedoninæ, charac. of, 670.
Eumedonius, 670.

- Eumenia crassa*, 131, 134.
Eumenidæ (Sandwich Is.), 678.
Eumolpidæ, new sp. & gen., by J. S. Baly, 246.
Eunoa nodosa, 129, 134; *Ærstedii*, 129, 134.
Eupetes nigricrissus, 632.
Eupleurodon, charac. of, 650.
Euplœa, 290, 299; *abjecta*, 300; *alcatheö*, 301; *amymone*, 300; *andamensis*, 300; *Angasii*, 299; *anthracina*, 301; *Batesii*, 299; *baudiniana*, 299; *Boisduvalii*, 300; *belinda*, 299; *Brenchleyi*, 301; *callitheö*, 292; *core*, 291, 301; *coreoides*, 301; *corinna*, 299; *Dalmanii*, 301; *deione*, 301; *diana*, 300; *distincta*, 299; *dolosa*, 300; *Donovani*, 299; *Doubledayi*, 301; *Duponchelii*, 301; *eleuthe*, 299; *Escholtzii*, 299; *eunice*, 292; *eurypon*, 301; *eyndhovii*, 301; *Felderi*, 300; *Frauenfeldi*, 300; *geyeri*, 301; *Godartii*, 301; *Guerini*, 300; *helcita*, 299; *Horsfieldii*, 300; *Hübneri*, 300; *janus*, 300; *Kirbyi*, 300; *Leachii*, 300; *Lewinii*, 299; *lorquini*, 300; *megæra*, 300; *Ménézi*, 301; *mitra*, 300; *montrouzieri*, 299; *morosa*, 301; *orope*, 299; *paykullei*, 301; *Perryi*, 299; *pietretii*, 301; *Pinwilli*, 301; *plexipus*, 291; *Poeeyi*, 301; *prosperina*, 300; *redtenbacheri*, 299; *Schlerzeri*, 300; *Schmeltzi*, 300; *Semper*, 296; *siamensis*, 301; *similis*, 291; *Staintonii*, 293; *Swainsonii*, 299; *torvina*, 301; *vermiculata*, 301; *vicina*, 301; *violetta*, 300; *Wallengrenii*, 300; *Whitmei*, 300.
Euprognatha, charac. of, 645.
Eurydemus grandis, 259; *Jansoni*, 258.
Eurynolambrus, remarks on, 669.
Eurynome, charac. of, 659.
Eurypodiidæ (*Stimps.*), 644.
Eurypodius, charac. of, 646.
Eurystomus crassirostris, 686; *pacificus*, 686.
Eustira (footnote), 538.
Eutrochus gemmatus, 600.
Eutropiichthys, 563, 578.
Exostoma, 568, 578.
Eylais, sp., 121.
 Feilden, Capt., insects coll. by, in high northern latitudes, 98.
 Fermentation and animal organisms, 1-94; apparatus used in exper., 10, 15; bacillus spores with heat, 79; Bastian's exper., 38, and with urine and liq. potass., 43, 45; conditions fa-
 vouring, 1; Cohn, Eidam, & Koch's exper. 73; exper. with superheated fluids, 50; germ-theory of disease, 83; heat as a promoter of, 5; in hay infusion, 57; in milk, 59; in urine, 53; introductory remarks on, 1; liquor potassæ appar. used in experiments, 10, 11; oxygen as a promoter of, 9; Pasteur's views on germs, 90; Potassæ as a promoter of, 12; Richardson on glandular theory of disease (ftnote), 88; Robert's exper. on, 28; Sanderson, Prof. Burdon, on germs, 90; signs of with superheated fluids, 53; species of organisms found, 55, 58; Tyndall and Pasteur's exper., 30, 70.
Fibrenus bullatus, 482, 483.
Fibularia, 441.
Filaria Bancrofti, 356, 359, 364; résumé of literature thereon, 365, 368; minutus, 304; sanguinis hominis, 360, 363, 364; sanguinolenta, 311, 363; Wuchereri, 361.
Filaria, 304; Cobbold's summary of diseases &c. connected therewith, 366, 667; devel. cannot progress far in host containing parent worm, 304; embryos do not all attain maturity, 308; embryo, its escape, 304; future history of filaria, 311; how to procure mosquitos containing filariæ, 306; large proport. of filariæ ingested by mosquito, 307; metamorphosis of embryo, 308; mosquito after feeding on, 305; mosquito as nurse of, 305.
Filariöse dermathemica, 362.
Filigrana, 720.
 Finches, note on Australian, 95.
 Fisheria (*Lockington*), 665.
 Fishes, geograph. distrib. of Indian freshwater, 534.
Flagellifera, a group of Radiolaria, 174, 175; charac. of, 176.
Flustra membranacea, 498.
Fluta (ftnote), 557.
Foraminifera compared with Stromatopora, 227.
Forficula sp. ? (*Java*), 135.
Formica cinerea, 290; *fusca*, 265, 271, 273, 274, 276, 278, 280, 283, 287, 288, 290; *sanguinea*, 290.
Formicidæ of Sandwich Is., 675.
Fredericella, 490, 497.
Fritillaria, arctic, 124; *furcata*, var., 125.
Fungus mycelium in boiled hay infusion, 58.
Gadus callarias, 692; *macrocephalus*,

- 689; *macrophthalmus* (fnote), 689; *morrhua*, 692; *punctatus*, 690.
Gagata, 563, 578.
Garra (fnote), 536; *alta* (fnote), 540; *ceylonensis* (fnote), 540; *malabarica* (fnote), 540.
Gastropoda, 422.
Gaza (n. gen.), 586; charac. of, 601; *dædala*, 602.
Geochelidon, 403.
Geoffroyius aruensis, 685.
Geographical distribution of Gulls and Terns (*Laridæ*), by H. Saunders, 390; of *Hippidea*, 314; of Indian Fresh-water Fishes.—Part III. Conclusion, by F. Day, 534; of some *Coleoptera*, 530; of *Radiolaria*, 172.
Geometræ of arctic reg., 114.
Geopelia humeralis, 688.
Germs, as to survival of, 48.
Gibelion (fnote), 537.
Glycera capitata, 127.
Glyptosternum, 562, 565, 578.
Gnathopogon (fnote), 538.
Gobio anisurus (fnote), 542; *barbus* (fnote), 537; *bicolor* (fnote), 542; *boga* (fnote), 542; *Bovianus* (fnote), 542; *canarensis* (fnote), 546; *Hamiltonii* (fnote), 542; *isurus* (fnote), 543; *limnophilus* (fnote), 543; *lissorhynchus* (fnote), 542; *malacostomus* (fnote), 542; *rienorhynchus* (fnote), 542.
Gobionichthys (fnote), 537.
Gobius, 561, 564, 578.
Gonatonotus, charac. of, 670.
Goniothorax, charac. of, 671.
Gonoproktopterus (fnote), 538.
Gonorhynchus bimaculatus (fnote), 540; *brachypterus* (fnote), 540; *brevis* (fnote), 543; *caudatus* (fnote), 540; *fimbriatus* (fnote), 543; *McClellandi* (fnote), 540; *macrosomus* (fnote), 543; *petrophilus* (fnote), 540; *rupeculus* (fnote), 540; *stenorhynchus* (fnote), 540.
Goura Albertisii, 688.
Graeculus subularis, 630.
Gregarina of Allman Bay, 125.
Gryllus sp. ? (Java), 135.
Gulls and Terns, Geograp. distrib. of, by H. Saunders, 390.
Gygis, 404, 406; *candida*, 401; *micro-rhyneha*, 404.
Gymnocorax senex, 629.
Gymnotus notopterus (fnote), 557.
Habits of Ants, Bees, and Wasps, Sir J. Lubbock on, 265, 607.
Hadena algida, 122; *Richardsonii*, 122; *septentrionis*, 122.
Halecyon Macleayi, 686; *sancta*, 686.
Halelminthidæ (arctic), 131.
Haliastur girrenera, 685.
Halicalyptra, 178.
Halimus, charac. of, 646.
Haliomma, 168, 177.
Haliommatida, 177.
Haliommatidium, 153, 170, 178.
Haliotis, 606.
Haliphormis, 178.
Haliplana, 403.
Hampala (fnote), 538.
Hanley, Mr. S., description of two new Shells, 580.
Haplochilus (fnote), 559, 561, 566, 578; *argenteus* (fnote), 535; *lineatus*, 536; *melastigma*, 535; *panchax*, 536, 568, 571; *rubrostigma*, 535.
Harmothoë imbricata, 129, 134.
Harpyopsis novæ-guinææ, 627.
Harrovia, charac. of, 671.
Hart, Mr., insects collected by, in Arctic regions, 98.
Hawkshaw, Mr. H., on action of Limpets (*Patella*) on Chalk at Dover, 406.
Helenus hesiformis, 482, 485.
Heliodiscus, 151, 152, 168, 170.
Heliophorus, 530.
Heliosphæra, 168, 179; *actinota*, 168; *inermis*, 168, and skeleton of, 149; *tenuissima*, 168.
Helix ericetorum, 700.
Hematopinus trichechi, 118.
Hemerobius lutarius, 731, 736.
Hemibarbus (fnote), 538.
Hemipholis microdiscus, 447, 448, 467, 480.
Hemiptera, descrip. of new, 482; of arctic reg., 118.
Hemus, charac. of, 654.
Henicophaps albifrons, 633.
Herbstia, charac. of, 654, 655.
Herbstiella, charac. of, 655.
Heterocera (arctic reg.), 112.
Heterocrypta, charac. of, 669.
Heteropora cervicornis, 724; *neozelandica*, 725; *pelliculata*, 724; relations to *Stromatopora*, 239; recent species of, 724.
Heteroporella, relations to *Stromatopora*, 239.
Hieracidea orientalis, 627.
Hippa, 323; *analogæ*, 324; *asiatica*, 325; *emerita*, 323; *talpoidea*, 324, and early stages and habits of, 315.
Hippidea, affinities of, 313; charac. of, 316; geographical distrib. of, 314; habits of, 315; introductory remarks

- on, 312; list of species of, 314; revision of, by E. J. Miers, 312.
- Histiastrium*, 176.
- Holes in Chalk made by Limpets, 409.
- Holocentropus dubius*, 732.
- Homaloptera, 536, 562, 566, 579; bilineata, 540; Brucei, 539, 568 (ftnote), 572; maculata, 540, 568 (ftnote), 572.
- Homœodera pygmæa, 530.
- Hoplia aurantiaca*, 135.
- Huenia*, charac. of, 648.
- Hyalopomatus Claparedii*, 127.
- Hyas, charac. of, 654.
- Hyastenus*, charac. of, 658.
- Hydrachnidæ (arctic), 121.
- Hydractinia*, 193, 194 (ftnote), 195, 235; compared with *Stromatopora*, 235, 236.
- Hydrobates erythropus*, 488; *regulus*, 482, 488.
- Hydrobatidæ, 488.
- Hydrochelidon*, 402; *hybrida*, 403; *leucoptera*, 403; (*Sterna*) *nigra*, 403.
- Hydrocorallinæ, relations to *Stromatopora*, 240.
- Hydrometra metator*, 482, 486.
- Hydrometridæ, 406.
- Hydropsyche instabilis*, 731, 736.
- Hydrozoa compared with *Stromatopora*, 233.
- Hymeniastrum*, 176.
- Hymenophysa* (ftnote), 539.
- Hymenoptera (arctic), 102, 106; from Sandwich Islands, 674.
- Hypselobarbus* (ftnote), 537, (ftnote) 538.
- Ichneumon erythromelas*, 106.
- Ichneumonidæ (arctic reg.), 106.
- Inachidæ, charac. of, 640, 642.
- Inachinæ and Subdiv. (*M.-Ed.* pt.), 644, 645, 646, 647.
- Inachoides*, charac. of, 646.
- Inachus*, charac. of, 645.
- Indian Freshwater Fishes, geograph. distrib. of, 534.
- Infusoria (arctic), 125.
- Insects from Java, report on, by Ch. O. Waterhouse, 134; of arctic regions, R. M'Lachlan on, 98; analysis of groups obtained in, 101; remarks on local sp. of, 106.
- Ischnodemus inambitiosus*, 482, 484.
- Isotoma Besselsii*, 119.
- Jansonius (n. gen.), 264; *alternatus*, 264.
- Japanese Echinoderms, Mr. W. P. Sladen on, 424; analysis of, 443.
- Newt, skull of, 718.
- Japanese Ophiuroidea, 445.
- —, relations of fauna, 445.
- Javan Insects, report on collection, by Ch. O. Waterhouse, 134.
- Jeffreys, Dr. J. Gwyn, note on position of genus *Sequenzia* among Gastropoda, 605.
- , on shells dredged by Capt. St. John in Korea Strait, 418.
- Jerdonia*, 539, 563, 566, 579; *maculata*, 555.
- Kellia pumila*, 421, 423.
- Korean Seas, Asteroidea and Echinoidea of, 424; Ophiuroidea of, 445; shells dredged in, 418.
- Labechia*, 192, 194, 234; compared with *Stromatopora*, 234; *conferta*, 234.
- Labeo*, 537, 561, 575, 579; *angra*, 542; *ariza*, 542; *bata*, 542; *boga*, 542; *boggut*, 542; *cæruleus*, 541; *calbasu*, 541; *diplostomus*, 542; *Dussumieri*, 541; *dyocheilus*, 542; *falcatus* (ftnote), 542; *fimbriatus*, 541; *gonius*, 541; *kawrus*, 542; *kontius*, 541; *microlepidotus* (ftnote), 541; *microphthalmus*, 542; *nandina*, 541; *nigrescens*, 541; *nigripinnis*, 542; *nukta*, 542; *pangusia*, 542; *porcellus*, 541; *potail*, 541; *Reynauldi* (ftnote), 541; *rohita*, 541; *sindensis*, 542; *Stoliczka*, 541; *velatus* (ftnote), 541.
- Labeobarbus* (ftnote), 538; *microlepis* (ftnote), 546; *mosal* (ftnote), 545.
- Labidomera imperialis*, 355.
- Labocheilus* (ftnote), 537.
- Lagisca rarispina*, 129, 134.
- Lahaina* (*Dana*), 658.
- Lambrus*, charac. of, 668, 672.
- Lamprodiscus*, 178.
- Lamprosoma amazonum*, 348; *armatum*, 350; *Batesi*, 347; *canaliculatum*, 349; *cuneatum*, 350; *cupricolle*, 348; *hypochryseum*, 347; *tricolor*, 346; *tridentatum*, 349.
- Largus humilis*, 483; *lentus*, 482, 483.
- Laria Rossii* (= ? *Dasychira grønlandica*), 112.
- Laridæ, geograph. distrib. of, 390; number species of, 391.
- Larridæ (Sandwich Is.), 676.
- Larus affinis*, 395, 397; *argentatus*, 395; *atricilla*, 399, 400; *Audouini*, 398; *Belcheri*, 396; *Bonapartii*, 399; *brachyrhynchus*, 396; *brunneicephalus*, 399; *Bulleri*, 398, 405; *cachinnans*, 395; *californicus*, 395, 396; *Cassini*, 402; *cirrhocephalus*, 398; *crassirostris*, 396; *dominicanus*, 397; *Frank-*

- lini, 399; fuliginosus, 392, 396; fuscus, 397; gelastes, 398; glaucescens, 395, 405; glaucodes, 399, 405; glaucus, 394; Hartlaubi, 398, 405; Heermanni, 396; Hemprichii, 400; ichthyaëtus, 399; leucopterus, 394, 395; leucophthalmus, 400; maculipennis, 399; marinus, 397; melanocephalus, 399; minutus, 399; modestus, 396; nigricans, 392; novæ-hollandiæ, 397, 405; occidentalis, 395; pacificus, 397; phæocephalus, 398, 405; philadelphiæ, 399; ridibundus, 399, 405; Saundersi, 399; serranus, 399; scopulinus, 397, 398, 405; Scoresbii, 397; zonorhynchus, 395, 396.
Lasæa rubra, 420, 423.
Lasius flavus, 267, 268, 273, 284; niger, 271, 276, 278, 280, 287, 289; queen of, 270.
Laubuca guttata (ftnote), 553.
 Lawes, Rev. W. G., Coll. Birds of New Guinea, 685.
Lebias (ftnote), 535.
Lepidocephalichthys, 539, 563, 566, 579.
Lepidonaxia (*Targioni-Tozzetti*), 671.
Lepidopa (synon.), 331.
Lepidops, 331; myops, 333; scutellata, 332; venusta, 332.
Lepidoptera of Arctic Regions, 102, 108; habits of, 103.
Leprotæ fulva, 250; *Lewisi*, 251.
Leptocephalichthys Berdmorei, 554; guntea, 554; thermalis, 554.
Leptocerus aterrimus, 733; bilineatus, 732; dissimilis, 733, 736; tineoides, 732.
Leptogenys insularis, 675.
Leptomithrax, charac. of, 656.
Leptomys cochlearis, 580; gravis, 580, 581.
Lepton sulcatulum, 420, 423.
Leptopinæ (*Stimps.*), 653.
Leptopodia, charac. of, 643.
Leptopodiidæ (*Dana*, pt.), 642.
Leptopodiinæ, charac. of, 642, 643.
Leptopsia, charac. of, 663.
Leptopus (*Lamk.* pt.), 654.
Lestridinæ, 391.
Lestris Hardyi, 392.
Lethæus lepidus, 482, 484; pallidiner-vis, 485.
Leucippe, charac. of, 649.
Leuciscus acinaces (ftnote), 553; Alfredianus (ftnote), 550; binotatus (ftnote), 546; branchiatus (ftnote), 551; chitul (ftnote), 544; cultellus (ftnote), 553; dandia (ftnote), 549; Dussumieri (ftnote), 553; Duvaucelii (ftnote), 548; dystomus (ftnote), 549; elingulatus (ftnote), 551; flavus (ftnote), 549; harengula (ftnote), 554; lateralis (ftnote), 549; lineolatus (ftnote), 552; malabaricus (ftnote), 549; margarodes (ftnote), 550; microcephalus (ftnote), 549; pellucidus (ftnote), 544; presbyter (ftnote), 549; rasbora (ftnote), 549; rubripes (ftnote), 551; salmoides (ftnote), 551; sulphureus (ftnote), 548; xanthogramme (ftnote), 549.
 Lewis, Dr. T. R., on Filariæ, 358, 360.
 Lewis and Cunningham's exper. on ferment organisms, 92.
Libidoclea, charac. of, 652.
Libinia, charac. of, 662.
Limnobates, 486.
Limnognon lotus, 482, 488; lubricus, 482, 488.
Limnophilus bipunctatus, 730; centralis, 732; griseus, 728, 730, 735; marmoratus, 729; rhombicus, 729; sparsus, 735; stigma, 728, 735; subcentralis, 729; vittatus, 732.
 Limpets acting on Chalk at Dover, J. C. Hawkshaw on, 406; rate of their excavations, 408.
Lineidæ, Arctic, 133.
Lineus fusca (arctic), 134.
 Linnaeus's Caddis-flies (*Phryganea*), H. D. J. Wallengren and R. M' Lachlan on, 726.
Liocassis, 562, 578.
Lipeurus, sp.?, on *Procellaria glacialis*, 119.
Lipura, sp.?, 119.
Lissa, charac. of, 663.
Lissorhynchus (ftnote), 536.
Litharachnium, 178; tentorium, skeleton of, 150.
Lithelida, 173, 176.
Lithelius, 154, 168, 176.
Lithobotrys, 178.
Lithocampe, 178.
Lithocircus, 179.
Lithocorythium, 178.
Lithocyclia, 176.
Litholophida, charac. of, 177.
Litholophus, 153, 177.
Lithomelissa, 158, 178.
Lithopera, 178.
Lithopoma imperator, 714.
Lithoptera, 153, 177.
Lithornithium, 178.
Lobivanelles miles, 688.
Loftusia, 227, (ftnote) 229.
Lophophæna, 178.
Lopnopus, 490.
 Lord-fish, 689, 690, 691.

- Loxorhynchus*, charac. of, 651.
Loxosomæ (arctic), 127, 130.
 Lubbock, Sir J., on anatomy of ants, 738; on habits of ants, bees, and wasps, 265, 607.
Luciobarbus (ftnote), 538.
Lumbricidæ (arctic), 133.
Lumbriconereis fragilis, 131, 134.
Lumbrinereidæ (arctic), 131.
Lycæna Aquilo, 111.
Lychnocanium, 177, 178; skeleton of, 150.
Lycosa glacialis, 120.
Lycosides, 120.
Lygæidæ, 483.
 M'Intosh, Dr. W. C., on Annelids of N. Polar Exped., 126; on remarkably branched *Syllis* dredged by H.M.S. 'Challenger,' 720.
 M'Lachlan, R., on Insecta coll. in late Arctic Exped. 78°-83° N. lat., 98.
 M'Lachlan, R., and Pastor H. D. J. Wallengren, on analysis of sp. Caddisflies (*Phryganea*) described by Linnaeus in his 'Fauna Suecica,' 726.
Macroceloma (= *Pericera*), charac. of, 665.
Macrocheira, charac. of, 647.
Macrochirichthys (ftnote), 538.
Macrones, 565, 578.
Macroplæa, 292.
Maia, charac. of, 655.
Maïidæ, 640, 653, 654, 655, 657, 659, 660, 661.
Mainea, charac., Fam., &c. of, 638, 640, 653.
Maïoid Crustacea, Edw. J. Miers on, 634.
Mallophaga (arctic region), 118.
Malurus albiscapulatus, 687.
Mamestra? *Feildeni*, 112.
 Manson, Dr. P., on develop. *Filaria sanguinis hominis* and Mosquito as nurse, 304; on *Filiaræ*, 363.
 Marenzeller, Dr. E., on Arctic Annelids, 127.
Margarita aspecta, 708; *carinata*, 697; *obscura*, 712; *poculosa*, 712; *polaris*, 701; *striata*, 601, 701; *umbilicalis*, 700.
Mastacembelus, 534, 561, 565, 578.
Mastigochirus, 321; *gracilis*, 322; *quadrilobatus*, 322.
Mastigopus (nom. præoc.), 321.
Mayoa (ftnote), 536.
Medusoid gonophores (arctic), 123.
Megachile diligens, 684.
Megalostomis placida, 341.
Megarasbora (ftnote), 538.
Megascelis basalis, 340; *femorata*, 339; *posticata*, 339.
Melania Limborgi, 580.
Melanopyrrhus Robertsoni, 633, 687.
Melinna cristata, 127.
Melolonthidæ of Java, 135.
Menæthus, charac. of, 649.
Menopon gonophaeum, var. ?, 119.
Merops ornatus, 686.
Mesorhœa, charac. of, 669.
Metoporphis, charac. of, 643.
Micippa, charac. of, 661.
Micippinæ, 660, 661.
Micippoides, charac. of, 658.
Micristius (ftnote), 535.
Micrococci, appearance in boiled fluids, 1; from hay infusion, 55.
Microœca flavovirescens, 630.
Microgaster, sp., 107.
Microglossum aterrimum, 685.
Micromugil (ftnote), 535.
Microphrys, charac. of, 664.
Micropisia (*Stimps.*), 654.
Microrhynchinæ, charac. of, 651.
Microrhynchus, 651.
Microvelia mimula, 482, 487.
Micrura fusca, 133.
 Miers, E. J., on Classification of *Maïoid Crustacea* or *Oxyrhyncha*, 634; revision of the *Hippidea*, 312.
 Millar, Dr. J., opinions on *Loftusia* and *Parkeria* (ftnote), 229.
Millepora, 233; *aleicornis* (ftnote), 195.
Milnia (*Stimps.*), 664.
Mimulus, charac. of, 649.
Misgurnis (ftnote), 539; *lateralis* (ftnote), 554.
Mithracinæ, charac. of, 666.
Mithraculus, charac. of, 667.
Mithrax, charac. of, 667.
 Mivart, Prof. St. G., Notes on recent researches on the *Radiolaria*, 136.
Mocosoæ, charac. of, 648.
Mola (ftnote), 537, 538; *Buchanani* (ftnote), 544.
Mollusca of the 'Challenger' expedition, by the Rev. R. Boog Watson, Pt. I., 506; Pt. II., 508; Pt. III., 586; Pt. IV., 692.
Monarcha melanopsis, 687.
Monocyrtida, charac. of, 177.
Monocyttaria, 173.
Monopterus, 557, 563, 579; *cinereus* (ftnote), 558; *helvolus* (ftnote), 558; *javanensis*, 558; *marmoratus* (ftnote), 558; *xanthognathus* (ftnote), 558.
Monozoa, 173.
Morara (ftnote), 538.
Morrhua macrocephala, 689, 690, 691; *vulgaris*, 689, 691.

- Morulus (fnote), 537.
 Mosquito after feeding on human blood, 305; as nurse of Filaria, Dr. P. Manson on, 304, 305.
 Moss, Dr. Edw. L., on the surface-fauna of arctic seas, 122.
 Mouhotia (n. gen.), 262; femorata, 262.
 Mrigala Buchananii (fnote), 543.
 Mugil, 578.
 Munia caniceps, 688.
 Murie, Dr. J., on skull of Echidna from Queensland, 413.
 — and Prof. H. A. Nicholson on Stromatopora and allies, 187.
 Muscidae Acalyptera, 118; of arctic reg., 117.
 Myrianida, 720.
 Myrmica ruginodis, 270.
 Myrmicidae (Sandwich Is.), 675.
 Mystacides atra, 731; bifasciata, 732; nigra, 732; quadrifasciata, 732.
 Mytilus carinatus, 421.
 Myxobrachia, 179; pluteus, nuclear vesicle of, 141; concretions of, 144.
 Myxodia ? sp. (arctic), 116.
 Nænia inca, 404.
 Nais proboscidea, propag. by divis., 720.
 Nandina (fnote), 537.
 Nandus, 562, 564, 578.
 Nangra, 563, 578.
 Nanomia (arctic), 123; cara, 125.
 Nasiterna keiensis, 628.
 Naucoridae, 489.
 Nautilus, 606.
 Naxia, charac. of, 658.
 Naxioides (A. M.-Edw.) (and fnote), 658.
 Næra cochlearis, 581.
 Nemacheilichthys, 539, 579; Rüppelli, 555.
 Nemacheilus, 539, 566, 579; Beavani, 556; botia, 555; butanensis, 556; cincticauda, 556; corica, 555; Denisonii, 556; Evezardi, 555; gracilis, 556; Guentheri, 555; ladacensis, 556; marmoratus, 556; monoceros, 555; montanus, 556; multifasciatus, 556; notostigma, 556; pavonaceus, 555; pulchellus, 555; rubidipinnis, 555; rupicola, 556; savona, 556; semiarmatus, 555; sinuatus, 555; spilopecterus, 556; Stoliczkae, 556; striatus, 556; triangularis, 556; turio, 556; zonalternans, 556; zonatus, 556.
 Nemausa, charac. of, 666.
 Nemertinea (arctic), 133.
 Neovelina (n. gen.), 487; Trailii, 482, 487.
 Nephthys longisetosa, 127.
 Nereidae (arctic), 131.
 Nereis pelagica, 127; prismatica, 131; zonata, 131, 134.
 Neureclipsis bimaculata, 729, 735; figurinensis, 735.
 Neuronina clathrata, 727; phelæonoides, 726; reticulata, 727; rufierus, 727, 734; striata, 728.
 New Guinea, birds of, 626, 685.
 Nicholson, Prof. H. A., and Dr. J. Murie on Stromatopora and allies, 187.
 Nirmus cingulatus, 118; phænotus, 119.
 Noctuae of arctic reg., 112; and larvae of, 114.
 North Polar expedition, Annelids of Dr. W. C. McIntosh on, 126.
 Note as to position of genus *Sequenzia* among Gastropoda, by Dr. J. Gwyn Jeffreys, 605.
 — on Australian Finches of genus *Poephila*, by Capt. W. E. Armit, 95.
 — on presence of *Tachyglossus* and *Ornithorhynchus* in N. and N.E. Queensland, by Capt. W. E. Armit, 411.
 Notes touching recent researches on the Radiolaria, by Prof. St. G. Mivart, 136.
 Nothria conchylega, 127.
 Notice of some shells dredged by Capt. St. John in Korea Strait, by Dr. J. Gwyn Jeffreys, 418.
 Notidobia ciliaris, 732.
 Notioxenus ferrugineus, 530.
 Notolopos, charac. of, 657.
 Notophthalmus viridescens, skull of, 717.
 Notopteridae, 559, 560.
 Notopterus, 561, 566, 579; bontianus (fnote), 557; Buchananii (fnote), 557; chitala, 557; hypselosoma (fnote), 557; kapirat, 557; lopis (fnote), 557; ornatus (fnote), 557; Pallasii (fnote), 557.
 Nucinella miliaris, 420; ovalis, 420, 423.
 Nudibranchs of Eastern Seas, C. Collingwood on, 737.
 Nullipora incrassata, 226; officinalis, 226.
 Nullipores, African, 226; compared with Stromatopora, 225.
 Nuria, 538, 563, 566, 579; daurica, 549, 568, 571.
 Nychia cirrosa, 122, 130, 134.
 Observations on the habits of Ants, Bees, and Wasps, by Sir John Lubbock, Pt. V., 265; Pt. VI., 607.

- Ocinopsis, charac. of, 645.
 Oenus pallidus, 255.
 Octonema (fntote), 536.
 Odynerus agilis, 681; congruus, 680; dubiosus, 681; localis, 678; maurus, 679; montanus, 680; rubritinctus, 679.
 Oecetis ochracea, 732.
 Oethra, charac. of, 669.
 Oikopleura (arctic), 124; rufescens, 125.
 Olyra, 578.
 Omalacantha (*Hale Streets*), 665.
 Ommatida, 157, 173, 174, 175; charac. of, 176.
 Ommatocampe, 177.
 Ommatospyris, 177.
 On a remarkably branched Syllis dredged by H.M.S. 'Challenger,' by Dr. W. C. McIntosh, 720.
 On Butterflies in coll. B.M. referred to gen. *Euplaea*, Fabr., by A. G. Butler, 290.
 On conditions favouring fermentation and appearance of Bacilli, Micrococci, and Torulae in boiled fluids, by Prof. H. C. Bastian, 1.
 On develop. *Filaria sanguinis hominis*, and on Mosquito as a nurse, by Dr. P. Manson, 304.
 On recent species of *Heteropora*, by G. Busk, 724.
 On the action of Limpets (*Patella*) in sinking pits and abrading the surface of the chalk at Dover, by J. C. Hawshaw, 407.
 On the anat. of Elk (*Alces malchis*), by Prof. M. Watson and Dr. A. H. Young, 371.
 On the Asteroidea and Echinoidea of the Korean Seas, by W. P. Sladen, 424.
 On the geograph. distrib. of the Gulls and Terns (*Laridae*), by H. Saunders, 390.
 On the life-history of *Filaria Bancrofti*, by Prof. T. S. Cobbold, 356.
 On the minute structure of Stromatopora and its allies, by Prof. H. A. Nicholson and Dr. J. Murie, 187.
 On the occurrence of *Morrhua macrocephala* at the mouth of the Thames, by Francis Day, 689.
 On the relations of Rhabdopleura, by Prof. G. J. Allman, 581.
 On some Ophiuroidea from the Korean seas, by Prof. P. M. Duncan, 445.
 On the structure and develop. of skull in Urodelous Amphibia, by Prof. W. K. Parker, 717.
 O'Neill, Dr., on Filariae, 362.
 Onychodactylus, skull of, 718.
 Onychopriion, 403.
 Ophiacantha Dallasii, 448, 471; indica, 473; stellata, 446, 473.
 Ophiacanthinae, 448.
 Ophiactis affinis, 448, 469, 481; Mulleri, 481; Reinhardtii, 481; sex-radia, 448, 469, 481, 446; simplex, 481.
 Ophiarachna, 447.
 Ophiarthrum, 447.
 Ophicardia (fntote), 557; Phayriana (fntote), 558.
 Ophichthys punctatus (fntote), 558.
 Ophioccephalus, 562, 565, 578; gachua, 568, (fntote) 558, 571; micropeltes, 567, 571; striatus, 570.
 Ophioenemis, 447.
 Ophiocoma, 447.
 Ophioglypha albida, 446; Forbesi, 448, 449, 479; nodosa, 446; sculpta, 448, 455, 479; sinensis, 446, 448, 460, 479; Sladeni, 448, 458, 479; striata, 448, 452, 479; Stuwitzii, 446, 479.
 Ophiolepididae, 448.
 Ophiolepis mirabilis, 447, 448, 460, 479.
 Ophiomusium, 451, 479.
 Ophionereidinae, 448.
 Ophionereis annulata, 480; crassispina, 480; dubia, 446, 464, var. sinensis, 448, 464; squamata, 480; variegata, 448, 462, 464, 480.
 Ophiopeza, 447.
 Ophioplocus, 447.
 Ophiopsammium, 447.
 Ophiothela isidicola, 477, 481; mirabilis, 477; tigris, 477; Verrilli, 477.
 Ophiotrix koreana, 448, 473; koreana, var. nov., 476; spiculata, 476; violacea, 476.
 Ophiothyreus Goei, 451.
 Ophiotrichidae, 448.
 Ophisternon (fntote), 557.
 Ophiuroidea of the Korean seas, 445.
 Oplousia, charac. of, 656.
 Opsaridium (fntote), 538.
 Opsarius (fntote), 538; acanthopterus (fntote), 551; anisocheilus (fntote), 551; bicirrhatus (fntote), 551; brachialis (fntote), 551; cirrhatus (fntote), 551; dualis (fntote), 551; fasciatus (fntote), 551; gracilis (fntote), 551; isocheilus (fntote), 550; latipinnatus (fntote), 551; maculatus (fntote), 551; malabaricus (fntote), 551; megastomus

- (fnote), 551; piscatorius (fnote), 550.
 Oregonia, charac. of, 646.
 Orias (fnote), 539.
 Oreinus, 536, 573, 579; guttatus (fnote), 540; Hodgsonii (fnote), 540; maculatus (fnote), 540; plagiosomus, 540; Richardsonii, 540; sinuatus, 540.
 Oribata Lucasii, 121; triangularis, 121; sp.?, 121.
 Oribatidæ (arctic reg.), 121.
 Oriolus striatus, 687.
 Ornithology of New Guinea, 626, 685.
 Ornithorhynchus, 411, 413.
 Osphromenus, 562, 565, 572, 574, 578.
 Osten-Sacken, Baron von, report on arctic Diptera, 116.
 Osteobrama (fnote), 538; cotio (fnote), 550; rapax (fnote), 550.
 Osteochilus, 537, 575, 579; cephalus, 543; chalybeatus, 543; Neilli, 543.
 Othonia, charac. of, 665.
 Othoninæ (*Dana*, *Stimps.*), 665.
 Oxygaster (fnote), 538.
 Oxyrhyncha, or Maioid Crustacea, E. J. Miers on, 634.
 Oxytelus alutaceifrons, 530.
 Pachnephorus Brethinghami, 256; Lewisi, 257; porosa, 257; torridus, 256.
 Pachycephala leucogastra, 631.
 Pachystomus (fnote), 538.
 Pachystroma, 214, 223; antiqua, 224; (Syringostroma) densa, 223.
 Pagophila, 400, 406; eburnea, 400.
 Palæacis, 232; compared with Stromatopora, 233.
 Pamera bilobata, 484; pagana, 482, 484.
 Panchax (fnote), 535; Buchanani (fnote), 536; cyanophthalmus (fnote), 535; Kuhlîi (fnote), 536; melanopterus (fnote), 536.
 Pangasius, 562, 578.
 Pangio (fnote), 539.
 Panopea plicata, 421; var. carinata, 421.
 Paradanio (fnote), 538; aurolineatus (fnote), 552; elegans (fnote), 552.
 Paradisea raggiana, 687.
 Paralaubuca (fnote), 538.
 Paramaya (*De Haan*), 666.
 Paramicippa, charac. of, 662.
 Paramithrax, charac. of, 655, 656.
 Parascela (n. gen.), charac., 252; cribrata (*Schauf.* type), 253.
 Parasites on arctic animals, 118.
 Parathoë, charac. of, 666.
 Parker, Prof. W. K., on skull of Urodolous Amphibia, 717.
 Parkeria, 194, 227, 228.
 Parthenope, charac. of, 668.
 Parthenopidæ, charac. of, 641, 667; and Subdiv., 668, 669, 670.
 Parthenopinæ, 641, 668, 669.
 Parthenopoides, charac. of, 672.
 Paryphes flavicollis, 483; (Sundarus) pontifex, 482; regalis, 483.
 Patella, action of, on chalk at Dover, J. C. Hawkshaw on, 406.
 Pecchiolia (or Verticordia) acuticostata, 420.
 Pecten similis, 419, 423.
 Pectinatella, 490, 497; magnifica, 505.
 Pectinura, 447.
 Pedicellariæ, Dr. W. P. Herapath's researches on, and terminology (fnote), 433.
 Pelecus affinis (fnote), 553; diffusus (fnote), 553.
 Pelia, charac. of, 659.
 Pelocoris binotulatus, 489; procurrens, 482, 489.
 Peloplastus (*Gerstæcker*), 654.
 Pelopæus flavipes, 676.
 Penthina, sp. ? (arctic), 116.
 Pericera, charac. of, 664.
 Periceridæ, charac. of, 640; and Sections of, 662, 663, 664, 665, 666.
 Pericerinæ, 662.
 Perichlamydium, 176.
 Peridinea of Melville Bay, 125.
 Peridineum acuminatum, 126; Michælias, 126.
 Perilampus, 538, 563, 566, 579; atpar, 552; canarensis (fnote), 552; ceylonensis, 553; fulvescens (fnote), 553; guttatus (fnote), 553; laubuca, 553; macropodus (fnote), 552; macrurus (fnote), 549; mysoricus (fnote), 552; osteographus (fnote), 552; perseus (fnote), 553; psilopteromus (fnote), 552; recurvirostris (fnote), 549; reticulatus (fnote), 552; striatus (fnote), 552; thermophilus (fnote), 549.
 Perinea (*Dana*), 664.
 Periophthalmus (fnote), 559, 561, 578.
 Perla, 731.
 Petalospyris, 178.
 Pheidole pusilla, 675.
 Phlogœnas jobiensis, 634.
 Phryganea (Caddis-flies), H. D. J. Walengen and R. M'Lachlan on, 726.
 Phryganea albifrons, 729, 732; azurea, 729, 732; bicaudata, 731; bilineata,

- 732; bimaculata, 729, 730, 735; ciliaris, 732; filosa, 732; flava, 732; flavilatera, 730, 736; fusca, 732; grandis, 727, 734; grisea, 728, 735; hectica, 732; longicornis, 729, 732; minuta, 732; nebulosa, 732; nigra, 731; phalænoides, 726; reticulata, 727; rhombica, 728, 729; saltatrix, 732; striata, 727, 734; umbrosa, 732; Wæneri, 732.
- Phycodes (*M.-Edw.*), charac. of, 656.
- Phylactolæmatous Polyzoa, development of, 497; endocyst, structure of, 491; formation of statoblast of, 502; new forms of, 505.
- Phyllidia spectabilis, 738.
- Phyllodoce grœnlandica, 130, 134; Luetkeni, 127.
- Phyllodocidæ, arctic, 130.
- Physematium, 138, 140, 141, 179.
- Phytophagous Coleoptera, new species of, J. Baly on, 336.
- Picrocerus, charac. of, 661.
- Pinacocystis rubicunda, 181.
- Pinarolestes megarhynchus, 630.
- Pisa, charac. of, 657.
- Pisoides, charac. of, 657.
- Pison hospes, 676; iridipennis, 676.
- Pitta novæ-hollandiæ, 687.
- Plagiacantha, 151, 179.
- Planaria fusca, 133.
- Planetis, 403.
- Planorbulina, 422.
- Platacanthus (fnote), 539.
- Platycara (fnote), 536; anisura (fnote), 540; australis (fnote), 539; lisso-rhynchus (fnote), 540; nasuta (fnote), 540; notata (fnote), 540.
- Platylambrus, remarks on, 671.
- Platytypes (*Lockington*), 667.
- Platyrhynchus (*Desb. & Schramm*), 653.
- Platyschisma, 593.
- Pleistacantha, charac. of, 646.
- Plethopora cervicornis, 724.
- Pleurobranchia (arctic), 123; rhododactyla, 125.
- Pleurophricus (*A. M.-Edw.*), 660.
- Pleurotomaria, 606.
- Ploem, Dr. J. C., insects coll. by, and described by Ch. O. Waterhouse, 134.
- Plusia parilis, 114; quadriplaga, 114.
- Pneumabanchus (fnote), 557; albinus (fnote), 558; leprosus (fnote), 558; striatus (fnote), 558.
- Podargus papuensis, 686.
- Podocheila, charac. of, 643.
- Podocyrtes, 178.
- Podonema, charac. of, 643.
- Podopisa (*Hilgendorf*), 658.
- Podura hyperborea, 119.
- Poëphila Gouldiæ and P. mirabilis, Capt. W. E. Armit on, 95.
- Polistes aurifer, 682.
- Polyacanthus, 562, 565, 572, 578; signatus, 568, 571.
- Polycentropus flavomaculatus, 732.
- Polycollida, 179.
- Polycyrtida, charac. of, 177.
- Polycystina, 174, 175; charac. of, 177.
- Polycyttaria, 173.
- Polyergus, 268; rufescens, 276.
- Polyonidæ (arctic), 127, 129.
- Polysphæra, 178.
- Polyzoa compared with Stromatopora, 237; Phylactolæmatous, 489.
- Ponera contracta, 675.
- Poneridæ (Sandwich Is.), 675.
- Popilia biguttata, 135.
- Porites discoidea, 191.
- Porphyrio melanopterus, 688.
- Potamorrhaphis (fnote), 534.
- Preliminary notice on surface-fauna of arctic seas, Dr. Edw. L. Moss on, 122.
- Prenelepis clandestina, 675.
- Priapulius caudatus, 127.
- Prionorhynchus, charac. of, 662.
- Prismatium, 179.
- Pristolepis, 562, 564, 567, 571, 578; fasciatus (fnote), 560.
- Proctotrypidæ (arctic reg.), 107.
- Proechidia, 417.
- Prosopis Blackburni, 682; facilis, 683; fuscipennis, 682; hilaris, 683; volatilis, 683.
- Prostheacanthus (fnote), 539; spectabilis (fnote), 554.
- Prosthetops (n. gen.), 533; capensis, 533.
- Psammechinus intermedius, 435.
- Pseudecheneis, 578.
- Pseudeutropius, 562, 565, 578.
- Pseudocolaspis, charac. of, 248; cærulea (*Laporte*, type), 248; femorata, 261; Lefevrei, 260; longicollis, 259.
- Pseudomicippe, charac. of, 661.
- Pseudorectes ferrugineus, 687.
- Psilorhynchus, 536, 563, 579; balitora, 540; variegatus (fnote), 540.
- Psychophora Sabini, 114.
- Pterocanium, 178.
- Pterocodon, 178.
- Pteropoda, 423.
- Pteropsarion (fnote), 538.

- Ptilonopus aurantiifrons*, 688; *coronulatus*, 688; *iozonus*, 688.
Ptilorhis Alberti, 630.
Ptychina biplicata, 421.
Ptycobarbus, 537, 579; *conirostris*, 541.
Pugettia, charac. of, 650.
Puncturella noachina, 422; var. *princeps*, 422.
Puntius (fnote), 538, 575; *chrysopoma* (fnote), 545; *Hamiltonii* (fnote), 547; *modestus* (fnote), 548; *perlee* (fnote), 547; *phutuni* (fnote), 548; *sophore* (fnote), 549.
Pylosphæra, 178.
Pyralidæ (arctic reg.), 115.
Pyrelia cadaverina?, 117.
Pyria, charac. of, 663.
Pyromaia (*Stimps.*), charac. of, 651.
Pyxinia, sp., 125.
Quedius fulgidus, 107.
Queensland, *Echidna* and *Ornithorhynchus* in, 411.
Rachinia, charac. of, 663.
Radiolaria, notes on recent researches on, by Prof. St. G. Mivart, 136.
Radiolaria, *alveoli* of, 141, 143; arctic seas, 125; bilateral mode of growth, 170; capsule of, 137, 140, 162, 167, 181; centrifugal spines of, 151; classification of, 173; colonies of, 138, 167; concretions of, 142; cool temperature necessary to, 161; crystalline bodies of, 143; distinctions from *Heliozoa*, 180; extracapsular bodies, 143, 166; extracapsular sarcodæ of, 143; extreme delicacy of, 157; distribution of, 170; fatty bodies of, 142; fission of, 162; flagellum of, 146; found at surface and deep sea, 171; geological distribution of, 172; growth, modes of, 168; Hertwig's opinion as to capsules, 162; impressionability of, 156; intracapsular sarcodæ of, 142; literature of, 183; locomotion of, 156; macrospores of, 165; manufacture silica, 158; microspores of, 165; motions of, 157; multicellular nature of, 183; multipolar growth, 168; nucleus of, 142, 182; nutrition of, 156; problematical bodies of extracapsule, 145; Prof. Hæckel's work on, 139; pseudopodia of, 137, 146; paralysis produced by, 158; relations to other organisms, 180; reproduction and growth of, 159; sarcodæ of, 137, 140, 141, 146; Sir C. Wyv. Thomson's remarks on (fnote), 137; skeleton, siliceous, of, 147, its gradual increasing complexity, 148; spore reproduction, 167; unicellular nature of, 182; yellow cells of, 137, 144, 182, hepatic nature of, 158; zoospores of, 161, 168; yellow cells of, supposed parasitic, 145; zooids of, 137, 139; zoospores with and without crystals, 162, 164, 165.
Rasbora, 538, (fnote) 559, 561, 566, 579; *Buchanani*, 549; *daniconius*, 549; *elanga*, 549; *neilgherriensis* (fnote), 549; *woolaree* (fnote), 549.
Reinwardtæna Reinwardti, 633.
Remipes, 316; *adactyla*, 318; *barbata*, 320; *cubensis*, 320; *ovalis*, 318; *pictus*, 318; *scutellatus*, 319; *strigillatus*, 320; *testudinarius*, 316; var. *denticulatifrons*, 318; *truncatifrons*, 321.
Report on Insecta (including *Arachnida*) of Arctic Expedition, by R. M'Lachlan, 98.
— — — obtained by Dr. J. C. Ploem in Java, with descrip. of new species of *Hoplia*, by Ch. O. Waterhouse, 134.
Revision of the *Hippidea*, by E. J. Miers, 312.
Rhabdopleura, 489; absence of endocyst and contractile cord of, 584; ectocyst of, 585; lophophore of, 581; *mirabilis*, 584, (fnote) 581; *Normani* (fnote), 581, 584; shield, import of, 584; tentacular cyst, 582.
Rhamphistoma (fnote), 534.
Rhaphidococcus, 152, 168, 177.
Rhaphidozoum, 179.
Rhipidura maculipectus, 631.
Rhitidoceros ruficollis, 686.
Rhizosphæra, 155, 168, 170.
Rhodeus indicus (fnote), 544; *macrocephalus* (fnote), 544.
Rhodia (*Bell*), 654.
Rhodostethia rosea, 400, 406.
Rhopalastrum, 176.
Rhopalocanium, 178.
Rhopalocera (arctic reg.), 108.
Rhyacophilidæ, 731.
Rhynchobdella, 578.
Rhynchopus albicollis, 405; *flavirostris*, 405; *nigra*, 405.
Rhynchopsinæ, geog. distrib. of, 390, 405.
Rhyparida sp. ? (Java), 135.
Richardson, Dr., on glandular theory of disease (fnote), 88.
Risella, 603.
Rissa brevirostris, 394; *tridactyla*, 392, 394.
Rita, 563, 578.

- Rohita (fnote), 537; Belangeri (fnote), 541; Buchanani (fnote), 541; Duvaucelii (fnote), 541; Leschenaultii (fnote), 541; lineata (fnote), 543; Reynauldi (fnote), 541; Rouxii (fnote), 541.
- Rohitichthys (fnote), 537.
- Rohtee, 538, 563, 566, 579; Bakeri, 550; Belangeri, 550; cotio, 550; Neilli, 550; Ogilbii, 550; Vigorsii, 550.
- Sabella sp.? in stomach of Phoca barbata, 127; spetsbergensis, 133, 134.
- Sabellidæ (arctic), 133.
- Saccobanchus, 563, 565, 574.
- Sagitta bipunctata?, 124.
- Sagra ferox, 337; longipes, 337.
- Salamandra maculosa, skull of, 717.
- Salisbury, Dr., on Entozoa, 357.
- Salmacis sulcata, 439, 443, 444.
- Salmophasia (fnote), 538; oblonga (fnote), 553.
- Salpinx (typical), 292; ægyptus, 295; alcidice, 294; assimilata, 293; callithoë, 292; chloë, 295; consanguinea, 293; crassa, 295; diocletianus, 294; eleusine, 293; elisa, 292; Erichsonii, 295; eunice, 292; eupator, 294; fraterna, 293; frigida, 293; gamella, 295; graeffiana, 293; Hewetsonii, 293; hisme, 292; Hobsonii, 293; hyacinthus, 294; illustris, 294; imitata, 293; iphianassa, 293; kadu, 293; klugii, 294; lorenzo, 294; Lowii, 294; margarita, 294; mniszchii, 293; Ochsenheimeri, 294; pasithea, 293; phænareta, 292; phœbus, 292; radamanthus, 294; Rogenhoferi, 294; semicirculus, 292; splendens, 294; superba, 294; treitschkii, 294; usipetes, 294; vestigiata, 293; viola, 293.
- Sandwich Islands, new Hymenoptera from, 674.
- Saunders, Howard, on geog. distrib. of Gulls and Terns (Laridæ), 390.
- Saxicava rugosa, 422, 423.
- Scacchia elliptica, 421.
- Scalibregma inflatum, 127.
- Scalibregmidæ (arctic), 131.
- Scaphiodon, 537, 566, 568, 572, 579; brevidorsalis, 544; irregularis, 544; Nashii, 544; Thomassi, 544; Watsoni, 544.
- Scaphisoma elongatum, 533.
- Scatophaga apicalis, 118.
- Scelodonta bidentata, 252; egregia, 252; Simoni, 251.
- Schacra (fnote), 538.
- Schismatorhynchus (fnote), 537.
- Schistura (fnote), 539; aculeata (fnote), 554; punctata (fnote), 555; subfusca (fnote), 556.
- Schizaster canaliferus, 442; ventricosus, 442, 443, 444.
- Schizophrys, charac. of, 660.
- Schizophrysinæ, 659.
- Schizopygopsis, 536, 579; Stoliczkae, 540.
- Schizothorax, 537, 578; esocinus, 540; progastus, 540.
- Sciæna, 578.
- Sciara ♂, sp.? (arctic reg.), 117.
- Sciæna lobata, 132, 134.
- Scirus, examples of, 121.
- Scissurella, 606.
- Scombresocidæ, 534, 559, 560.
- Scoparia gelida, 115.
- Scrobicularia adunca, 581.
- Scudder's views of Euplœa, 291.
- Scyra, charac. of, 663.
- Seythrops novæ-hollandiæ, 628, 686.
- Seгуenzia, 586; carinata, 590, 606; elegans, 606; formosa, 587, 605, var. lineata, 587; ionica, 589; trispinosa, 591.
- Seironota perspicillata, skull of, 717.
- Semiplotus, 537, 563, 579; M'Clelandi, 543; modestus, 543.
- Sharpe, R. Bowdler, on birds of New Guinea, Pt. V., 626; Pt. VI., 685.
- Shells, descriptions of two new, 580; of 'Challenger' exped. 508, 586, 692; of Korean seas, 418.
- Sialis lutaria, 730, 736.
- Sicydium, 562, 564, 578.
- Siliquaria, 606.
- Silundia, 563, 578.
- Siluridæ, 559, 560.
- Silurus, 565, 568, 578.
- Silva Lima, Dr., on Filariæ, 362.
- Simocarcinus, charac. of, 649.
- Siphodontalium bifissum, 521; dicellum, 521; eboracense, 523; filum, 520; gracile, 520; lofotense, 520; minutum, 520; pentagonum, 520; platamodes, 519; prionotum, 522; pusillum, 520; quinquangulare, 519; tetragonium, 516; tetraschistum, 521; tythum, 520; vitreum, 520.
- Siphonosphæra, 168, 179.
- Sisor, 563, 578.
- Skull of Urodelous Amphibia, Prof. W. K. Parker on, 717.
- Sladen, W. P., on Asteroidea and Echinoidea of Korean seas, 424.
- Smiliogaster (fnote), 538.
- Smith, Fred., on Aculeate Hymenoptera from Sandwich Islands, 674.

- Solariella aureonitens*, 595; *trochoides*, 600.
Solarium, 605; *hybridum*, 606.
Solenolambrus, charac. of, 668.
Solenopsis geminata, 676.
Somileptes, 539, 563, 579; *bispinosa* (fthote), 554; *gongota*, 554; *unispina* (fthote), 555.
 Sonsino, Dr., on *Filariæ*, 361.
Sparsispongia, 190, 229.
Speckled Cod, 689.
Sperlepes rubra, skull of, 718; *salmonæa*, skull of, 718.
Sphærechinus, 434.
Sphæroidea, charac. of, 179.
Sphærospongia, 229.
Sphærozoa, 166.
Sphærozoida, 173; charac. of, 179.
Sphærozoum, 138, 159, 168, 172, 179; *ovo-di-mare*, skeleton of, 148.
Sphecotheres Salvadorii, 687.
Sphegidæ (Sandwich Is.), 676.
Sphenocarcinus, charac. of, 663.
Spirillina, 178.
Spirorbis lucidus, 127.
Sponges compared with Stromatopora, 229.
Spongia stellata, 229.
Spongida (Radiolaria), charac. of, 176.
Spongoastericus, 176.
Spongocyelia, 154, 176.
Spongodictyum, 155, 168.
Spongodiscida, 175.
Spongodiscus, 154, 176.
Spongosphæra, 155, 168.
Spongosphærida, 175, 176.
Spongotrochus, 176.
Spongurida, 173; an unnatural group, 175.
Spongurus, 154, 170, 176.
Spyridobotrys, 178.
Staurolithium, 153, 177.
Stellaster Belcheri, 430, 433.
Stellispongia, 229.
Stenocionopinæ, 652.
Stenocionops, charac. of, 652.
Stenophylax, 730.
Stenorhynchus, charac. of, 643.
Stephanastrum, 154, 176.
Stereorariinæ, geogr. distrib. of, 390, 391.
Stereorarius antarcticus, 393, 394; *asiaticus*, 392; *catarrhactes*, 392, 393, 394; *chilensis*, 393, 394; *crepidatus*, 391, 392; *parasiticus*, 391, 392; *pomatorhinus*, 392.
Sterna albigena, 402; *aleutica*, 403, 405; *anæstheta*, 404, 405; *anglica*, 403; *antarctica*, 402, 406; *balænarum*, 403; *Bergii*, 402; *cantiaca*, 401, 402; *caspia*, 401; *Dougalli*, 403; *elegans*, 401; *eurygnathus*, 401; *fluvialilis*, 401; *frontalis*, 402; *fuliginosa*, 404, 405; *hirundinacea*, 402, 406; *lunata*, 404, 405; *macrura*, 401; *magnirostris*, 403; *maxima*, 401; *media*, 401; *melanauchen*, 403; *minuta*, 403; *nereis*, 403; *nigra*, 403; *seena*, 403; *trudeaui*, 403; *virgata*, 402.
Sterninæ, geogr. distrib. of, 390, 401.
Steropsocus immaculatus, 733.
Stichocyrtida, charac. of, 178.
Stictoplea, 301; *binotata*, 303; *doleschalii*, 303; *euphon*, 303; *gloriosa*, 301; *grotei*, 302; *harrisii*, 302; *hopei*, 302; *immaculata*, 303; *inconspicua* ♂, 302; *inæqualis*, 302; *lankana*, 302; *lætifica*, 302; *microsticta*, 302; *mæsta*, 303; *palla*, 303; *pelor*, 303; *picina*, 302; *schlegelii*, 301; *superba*, 301; *swinhoei*, 302; *sylvester*, 303; *tristis*, 303.
Stilbognathus, charac. of, 652.
Stromatocerium, 189, 210, 222; *canadense*, 223; *rugosum*, 213, 223.
Stromatopora, minute structure of, and allies, by Prof. H. A. Nicholson and Dr. Murie, 187.
 —, affinities, 224; *calcareous*, 196, 231; *classification and types*, 216; *comparisons with Nullipores*, 225; — *Foraminifera*, 227; — *corals*, 232; — *Hydrozoa*, 233; *concentric arrang. chamb. round vert. column*, 212; *departures from type*, 212; *fossilization*, 196; *gen. struct. of typical Stromatopora*, 195; *hexactinellid nature of*, 230; *history and literature*, 188; *horiz. laminae*, 205; *horny*, 197, 229; *internal constitution*, 203; *introductory remarks*, 187; *minute structure*, 202; *mode of occurrence*, 196; *nature of radial pillars*, 203; *original constitution*, 196; *radiated water-canals*, 209; *sarcoide chambers*, 208; *siliceous*, 196, 230; *spongiöse nature of*, 231; *summary and conclusions*, 239; *systematic position*, 224; *tubes of *Caenopora**, 211; *vertical water-canals of*, 210.
Stromatopora, 217; *astroites*, 210; *concentrica*, 188, 189, 193, 195, 230; *constellata*, 210; (*Cænostroma*) *discoideum*, 218, 219, 233; *elegans*, 210; *expansa*, 218; *granulata*, 205, 210, 218, 219; *Hindei*, 209, 210; (*C.*) *incrustans*, 210; *nodulata*, 206, 207, 218; *nulliporoides*, 218, 219; *ostio-*

- lata, 210, 218, 219; (Caunopora) perforata, 218; polymorpha, 188, 189, 190, 217, 218, 230; ponderosa, 210; rugosa, 195; Schmidtii, 210; solidula, 210; striatella, 205, 210, 215; subtilis, 189; tuberculata, 204, 205, 210, 218, 233, 238; typica, 210, 218.
- Stromatoporoids, 189, 194, 198, 210, 213, 214, 215, 216, 237.
- Strongylocentrotus intermedius, 434, 443, 444.
- Stylocyclia, 176.
- Stylodictya, 168, 176.
- Stylodictyon, 221; (Syringostroma) columnæ, 221; retiforme, 212, 222.
- Stylospira, 154, 176.
- Stylospongia, 175, 176.
- Syagrus rubifrons, 263.
- Syllis fasciata, 127; ramosa, 720; remarkably branched form, Dr. W. C. McIntosh on, 720.
- Syma torotoro, 686.
- Symbranchidæ, 559, 560.
- Symbranchus, 557, 563, 566, 579; bengalensis, 558; eurychasma (fnote), 556; grammicus (fnote), 558; immaculatus (fnote), 558.
- Syncrossus (fnote), 539.
- Synopsis of Maioid Crustacea or Oxyrhyncha, E. J. Miers on, 634, 642.
- Syringostroma, 192, 193; columnaris, 210, 212; densa, 206, 207, 210.
- Systomus (fnote), 538; apogonoides (fnote), 547; assimilis (fnote), 549; carnaticus (fnote), 547; chrysostomus (fnote), 545; conchonioides (fnote), 548; gibbosus (fnote), 548; immaculatus (fnote), 545; leptosomus (fnote), 548; macularis (fnote), 547; madraspatensis (fnote), 549; malacopterus (fnote), 548; microlepis (fnote), 550; pyrrhopterus (fnote), 548; rubroinctus (fnote), 547; sophore (fnote), 548; tripunctatus (fnote), 548; tristis (fnote), 547.
- Tachina hirta, 117.
- Tachinidæ (arctic reg.), 117.
- Tachyglossus, 411; histrix, 413, 415; Lawesii, 413.
- Tachypetes minor, 688.
- Tadorna radjah, 688.
- Talegallus fuscirostris, 634.
- Tanyptera microrhyncha, 629, 686; sylvia, 629.
- Tarentula exasperans, 120.
- Taricha torosa, skull of, 718.
- Tegenaria detestabilis, 120.
- Teleophrys (*Stimps.*), 667.
- Tellimya ovalis, 422.
- Telmessus (*White*), remarks on, 670.
- Temnechinus, 438; maculatus, 438.
- Temnnotus, charac. of, 659.
- Temnopleurus Hardwickii, 435, 443, 444; Reynaudi, 437; toreumaticus, 438, 443, 444.
- Terebellidæ (arctic), 132.
- Terebellides Stroemi, 127.
- Terebratula caput-serpentis, var. septentrionalis, 418, 423.
- Terns and Gulls, geograph. distrib. of, by H. Saunders, 390.
- Tetrabanchus (fnote), 557; microphthalmus (fnote), 558.
- Tetramorium guineense, 675.
- Tetrapyle, 168, 177.
- Thalassicolla, 138, 139, 141, 159, 160, 161, 172, 179; nucleata, 156.
- Thalassolampe, 140, 179.
- Thalassoplaneta, 160.
- Thelopus circinnatus, 132, 134.
- Therediids, 120.
- Thoë, charac. of, 667.
- Thynnichthys, 537, 563, 571, 579; cochinchinensis (fnote), 544; sandkhol, 544.
- Thyrsocyrtis, 178.
- Tiarinia, charac. of, 664.
- Tinea Robertella, 733, 736.
- Tingitidæ, 485.
- Tinodes luridus, 732.
- Tinoporus as compared with Stromatopora, 227.
- Tintinnus (arctic), 126.
- Tipula arctica, 116, 117.
- Tortricidæ (arctic reg.), 116.
- Torulæ corpuscles, 58; from hay infusion, 55; their appearance in boiled fluids, 1.
- Traquaria, on position of, 172.
- Trematodiscida, charac. of, 176.
- Trematodiscus, 153, 154, 176.
- Trepischrois, 296; claudia, 296; midamus, 297; mulciber, 296; tisiphone, 297.
- Trevelyanella felis, 738.
- Trichina cystica, 357; spiralis, 364.
- Trichocera relegationis, 117.
- Trichogaster, 563, 572, 574, 578.
- Trichoglossus massena, 686.
- Trigonothir, charac. of, 648.
- Triopa principis, 738.
- Trochidæ, 586.
- Trochus Adansoni, 423, var. affinis, 702; (Margarita) ægleës, 704; amabilis, 601, 704; aureonitens, 595; (M.) azorensis, 710; (M.) brychius, 699; (M.) charopus, 703, and var. cæruleus, 701; cinereus, 601, 704;

- (M.) *clavatus*, 705; (Thalotia) *elisa*, 699; (M.) *gemma*losa, 709; (Gibbula) *glyptus*, 694; (M.) *groenlandica*, 711; *helcinus*, 713; (M.) *infundibulum*, 707; *iris*, 605; (M.) *lima*, 703; *majus*, 695; (M.) *nitens*, 713; *occidentalis*, 696; (M.) *pachychiles*, 708; (M.) *pompholugotus*, 702, 711; (M.) *rhysus*, 706; (M.?) *scintillans*, 712; (*Ziziphinus*) *strophorus*, 695; (Z.) *tiara*, 696, 705; *transenna*, 696; *tumidus*, 711; *turbinoides*, 423.
- Tropidorhynchus novæ-guineæ*, 687.
- Turbo*, 606; *coccineus*, 423; (*Calcar*) *henicus*, 713; (*Collonia*) *indutus*, 715; *philippiana*, 714; *purpureus*, 423; (*Guildfordia*) *rhodostoma*, 714; *sanguineus*, 423; *transenna*, 714.
- Turcica monolifera*, 708.
- Tyche*, charac. of, 653.
- Tylocarcinus*, charac. of, 664.
- Tylognathus* (ftnote), 537; *barbatulus* (ftnote), 543; *striolatus* (ftnote), 542; *valenciennesii* (ftnote), 542.
- Tylosurus* (ftnote), 534.
- Typica* (subsect. *Radiolaria*), charac. of, 177.
- Unibranchapertura* (ftnote), 557; *lævis* (ftnote), 558.
- Urodelous Amphibia*, struct. and devel. skull of, 717.
- Varicorhinus* (ftnote), 538; *bobree* (ftnote), 541.
- Velia virgata*, 482, 486; *vivida*, 482, 486.
- Veliidæ*, 486.
- Vesiculata*, a group of *Radiolaria*, 174, 175; charac. of, 179.
- Vespidæ* (Sandwich Is.), 682.
- Vibrio Rugula* from hay infusion, 55, 59; *serpens* (ftnote), 55.
- Wallago*, 562, 565, 578.
- Wallengren, Pastor H. D. J., and R. M'Lachlan, on Caddis-flies (*Phryganea*) of Linnæus's 'Fauna Suecica,' 726.
- Walrus parasite, 118.
- Wasps and bees, Sir J. Lubbock on, 265, 607, 625; conduct towards friends, 626; exper. as to knowledge of colour, 625.
- , new species from Sandwich Islands, 682.
- Waterhouse, Ch. O., report on insects from Java, coll. by Dr. J. C. Ploem, 134.
- Waterhouse, F. H., descriptions of new Coleoptera, 530.
- Watson, Prof. M., and Dr. A. H. Young, on anat. of Elk, 371.
- Watson, Rev. R. Boog, Mollusca of 'Challenger' Expedition, Pt. I., 506; Pt. II., 508; Pt. III., 586; Pt. IV., 692.
- White, Dr. J. Buchanan, descrip. of new Hemiptera, 482.
- Wucherer, Dr. O., on *Filariæ*, 357.
- Xema furcata*, 400; *Sabinii*, 400.
- Xenocarcinus*, charac. of, 648.
- Xiphacantha*, 177; sp. (*Wyr. Thoms.*), 152, 153.
- Xylocopa æneipennis*, 684.
- Young, Dr. A. H., and Prof. M. Watson, on anat. of Elk, 371.
- Zebrida*, charac. of, 670.
- Zonites cellarius*, 594.
- Zygocyrtida*, charac. of, 178.
- Zygonectes* (ftnote), 535.
- Zygostephanus*, 151, 179; skeleton of, 150.

END OF THE FOURTEENTH VOLUME.

PRINTED BY TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.